arXiv:0806.1844v1 [hep-th] 11 Jun 2008

Two-field cosmological models and large-scale cosmic magnetic fields

Alexander A. Andrianov

Departament d'Estructura i Constituents de la Materia and Institut de Ciències del Cosmos (ICCUB),
Universitat de Barcelona, 08028, Barcelona, Spain
V.A. Fock Department of Theoretical Physics, Saint Petersburg State University, 198504, S. Petersburg, Russia

Francesco Cannata

Istituto Nazionale di Fisica Nucleare, Via Irnerio 46, 40126 Bologna, Italy

Alexander Y. Kamenshchik

Dipartimento di Fisica and Istituto Nazionale di Fisica Nucleare, Via Irnerio 46, 40126 Bologna, Italy
L.D. Landau Institute for Theoretical Physics of the Russian Academy of Sciences, Kosygin str. 2, 119334 Moscow, Russia

Daniele Regoli

Dipartimento di Fisica and Istituto Nazionale di Fisica Nucleare, Via Irnerio 46, 40126 Bologna, Italy

E-mail: `andrianov@bo.infn.it`, `cannata@bo.infn.it`, `kamenshchik@bo.infn.it` and `regoli@bo.infn.it`

Abstract. We consider two different toy cosmological models based on two fields (one normal scalar and one phantom) realizing the same evolution of the Bang-to-Rip type. One of the fields (pseudoscalar) interacts with the magnetic field breaking the conformal invariance of the latter. The effects of the amplification of cosmic magnetic fields are studied and it is shown that the presence of such effects can discriminate between different cosmological models realizing the same global evolution of the universe.

PACS numbers: 98.80.Cq, 98.80.Jk

Keywords: phantom dark energy, scalar fields, magnetic fields

1. Introduction

The discovery of the cosmic acceleration [1] and the search for dark energy responsible for its origin [2] have stimulated the study of different field models driving the cosmological evolution. Such a study usually is called the potential reconstruction [3], because the most typical examples of these models are those with a scalar field, whose potential should be found to provide a given dynamics of the universe. In the flat Friedmann models with a single scalar field, the form of the potential and the time dependence of the scalar field are uniquely determined by the evolution of the Hubble variable (up to a shift of the scalar field). During last years the models with two scalar fields have also become very popular. This is connected with the fact that some observations point out that the relation between the pressure and the energy density could be less than -1 [4]. Such equation of state arises if the matter is represented by a scalar field with a negative kinetic term. This field is called "phantom" [5]. Moreover, according to some observations [6] the universe undergoes a transition between normal and phantom phase. Such an effect is dubbed "phantom divide line crossing" [7]. In principle, the effect of phantom divide line crossing can be explained in the model with the only scalar field provided a special form of the potential and initial conditions is chosen [8] or in models with a non-minimally coupled scalar field [9]. However, the models with two scalar fields, one standard and one phantom, look more "natural" for the description of the phantom divide line crossing [10, 11, 12]‡.

In our preceding paper [12] we have studied the procedure of reconstruction of the potential in two-field models. It was shown that there exists a huge variety of potentials and time dependences of the fields realizing the same cosmological evolution. Some concrete examples were considered, corresponding to the evolution beginning with the standard Big Bang singularity and ending in the Big Rip singularity [13].

One can ask oneself: what is the sense of studying different potentials and scalar field dynamics if they imply the same cosmological evolution? The point is that the scalar and phantom field can interact with other fields and influence not only the global cosmological evolution but also other observable quantities.

One of the possible effects of the presence of normal and phantom fields could be their influence on the dynamics of cosmic magnetic fields. The problem of the origin and of possible amplification of cosmic magnetic fields is widely discussed in the literature [15]. In particular, the origin of such fields can be attributed to primordial quantum fluctuations [16] and their further evolution can be influenced by hypothetic interaction with pseudoscalar fields breaking the conformal invariance of the electromagnetic field [17, 18]. In the present paper we consider the evolution of magnetic fields created as a result of quantum fluctuations, undergoing the inflationary period with unbroken conformal invariance and beginning the interaction with pseudoscalar or pseudophantom fields after exiting the inflation and entering the Big Bang expansion stage, which is a part of the Bang-to-Rip scenario described in the preceding paper [12]. We shall use different field realizations of this scenario and shall see how the dynamics of the field with negative parity influences the dynamics of cosmic magnetic fields. To our knowledge the possible influence of the two-field dynamics, (when one of two (pseudo)scalar fields is a phantom one) on the magnetic fields was not yet discussed in the literature.

Speaking of cosmic magentic fields we mean the large-scale galactic, intergalactic or super-cluster magnetic fields of order from $10^{-6}G$ to $10^{-11}G$ with correlation from 100 kpc to several Mpc to the extent that they are originated from scalar and, possibly gauge field fluctuations after exiting the inflation. Their seeds may well have $10^{-18} - 10^{-27}G$ or less (see [15]).

---

‡ These two fields may have their origin in spontaneous breaking of primordial symmetries of moduli space with complex potential [11].

The structure of the paper is as follows: in Sec. 2 we recall the Bang-to-Rip scenario and describe some examples of different dynamics of scalar and phantom fields; in Sec. 3 we introduce the interaction of the fields (phantom or normal) with an electromagnetic field and write down the corresponding equations of motion; in Sec. 4 we describe the numerical simulations of the evolution of magnetic fields and present the results of these simulations; Sec. 5 is devoted to concluding remarks.

2. Cosmological evolution and (pseudo)-scalar fields

We shall consider a spatially flat Friedmann universe with the metric

$$ds^2 = dt^2 - a^2(t)dl^2. \tag{1}$$

Here the spatial distance element dl refers to the so called comoving coordinates. The physical distance is obtained by multiplying dl by the cosmological radius $a(t)$. We would like to consider the cosmological evolution characterized by the following time dependence of the Hubble variable $h(t) \equiv \dot{a}/a$, where "dot" denotes the differentiation with respect to the cosmic time t:

$$h(t) = \frac{t_R}{3t(t_R - t)}. \tag{2}$$

This scenario could be called "Bang-to-Rip" [12] because it is obvious that at small values of t the universe expands according to power law: $a \sim t^{1/3}$ while at $t \to t_R$ the Hubble variable explodes and one encounters the typical Big Rip type singularity. (The factor one third in (2) was chosen for calculation simplicity). In our preceding paper [12]) we considered a class of two-field cosmological models, where one field was a standard scalar field ϕ, while the other was a phantom one ξ. The energy density of the system of these two interacting fields is

$$\varepsilon = \frac{1}{2}\dot{\phi}^2 - \frac{1}{2}\dot{\xi}^2 + V(\phi,\xi). \tag{3}$$

Analyzing the Friedmann equation §

$$h^2 = \varepsilon, \tag{4}$$

we have shown that in contrast to models with one scalar field, there is huge variety of potentials $V(\phi,\xi)$ realizing a given evolution. Moreover, besides the freedom in the choice of the potential, one can choose different dynamics of the fields $\phi(t)$ and $\xi(t)$ realizing the given evolution. We have studied in [12] some particular exactly solvable examples of forms of potentials and time dependences of fields ϕ and ξ providing the evolution (2). Here we shall present and apply some of them. Consider the potential ‖

$$V_I(\xi,\phi) = \frac{2}{9t_R^2}\cosh^6(-3\phi/4)\exp(3\sqrt{2}\xi). \tag{5}$$

---

§ We use the following system of units $\hbar = 1, c = 1$ and $8\pi G = 3$. In this system the Planck mass m_P, the Planck length l_P and the Planck time t_P are equal to 1. Then when we need to make the transition to the "normal", say, cgs units, we should simply express the Planck units in terms of the cgs units. In all that follows we tacitly assume that all our units are normalized by the proper Planck units. Thus, the scalar field entering as an argument into the dimensionless expressions should be divided by the factor $\sqrt{m_P/t_p}$.

‖ The expression for the potential should be multiplied by the factor $m_P/(cl_P)$.

and the fields

$$\phi(t) = -\frac{4}{3}\text{arctanh}\sqrt{\frac{t_R - t}{t_R}}, \tag{6}$$

$$\xi(t) = \frac{\sqrt{2}}{3}\ln\frac{t}{t_R - t}. \tag{7}$$

(Here the expressions for the fields $\phi(t)$ and $\xi(t)$ should be multiplied by $\sqrt{m_P/t_P}$. For the relation between Planck units and cgs ones see e.g. [19] If we would like to substitute one of these two fields by the pseudoscalar field, conserving the correct parity of the potential, we can choose only the field ϕ because the potential V_I is even with respect to ϕ, but not with respect to ξ. In what follows we shall call the model with the potential (5), the pseudoscalar field (6) and the scalar phantom (7) "model I".

Consider another potential

$$V_{II}(\xi,\phi) = \frac{2}{9t_R^2}\sinh^2(3\xi/4)\cosh^2(3\xi/4)\exp(-3\sqrt{2}\phi). \tag{8}$$

with the fields

$$\phi(t) = \frac{\sqrt{2}}{3}\ln\frac{t}{t_R - t}, \tag{9}$$

$$\xi(t) = \frac{4}{3}\text{arctanh}\sqrt{\frac{t}{t_R}}. \tag{10}$$

This potential is even with respect to the field ξ. Hence our model II is based on the potential (8), the pseudophantom field (10) and the scalar field (9). They will be the fields with the negative parity which couple to the magnetic field.

3. Post-inflationary evolution of a magnetic field interacting with a pseudo-scalar or pseudo-phantom fields

The action of an electromagnetic field interacting with a pseudoscalar or pseudophantom field ϕ is

$$S = -\frac{1}{4}\int d^4x\sqrt{-g}(F_{\mu\nu}F^{\mu\nu} + \alpha\phi F_{\mu\nu}\tilde{F}^{\mu\nu}), \tag{11}$$

where α is an interaction constant and the dual electromagnetic tensor $\tilde{F}^{\mu\nu}$ is defined as

$$\tilde{F}^{\mu\nu} \equiv \frac{1}{2}E^{\mu\nu\rho\sigma}F_{\rho\sigma}, \tag{12}$$

where

$$E_{\mu\nu\rho\sigma} \equiv \sqrt{-g}\ \epsilon_{\mu\nu\rho\sigma},\quad E^{\mu\nu\rho\sigma} \equiv -\frac{1}{\sqrt{-g}}\ \epsilon^{\mu\nu\rho\sigma}, \tag{13}$$

with the standard Levi-Civita symbol

$$\epsilon_{\mu\nu\rho\sigma} = \epsilon_{[\mu\nu\rho\sigma]},\ \ \epsilon_{0123} = +1. \tag{14}$$

Variating the action (11) with respect to the field A_μ we obtain the field equations

$$\nabla_\mu F^{\mu\nu} = -\alpha\partial_\mu\phi\tilde{F}^{\mu\nu}, \tag{15}$$

$$\nabla_\mu\tilde{F}^{\mu\nu} = 0. \tag{16}$$

The Klein-Gordon equation for the pseudoscalar field is

$$\nabla^{\mu}\nabla_{\mu}\phi + \frac{\partial V}{\partial \phi} = -\alpha F_{\mu\nu}\tilde{F}^{\mu\nu}. \tag{17}$$

The Klein-Gordon equation for the pseudophantom field (which is the one that couples with the magnetic field in the model II) differs from equation (17) by change of sign in front of the kinetic term. In what follows we shall neglect the influence of magnetic fields on the cosmological evolution, i.e. we will discard the electromagnetic coupling in equation (17).

If one wants to rewrite these formulae in terms of the three-dimensional quantities (i.e. the electric and magnetic fields) one can find the expression of the electromagnetic tensor in a generic curved background, starting from a locally flat reference frame — in which it is well known the relation between electromagnetic fields and F — and using a coordinate transformation. It is easy to see that we have, for the metric (1):

$$F^{\mu\nu} = \frac{1}{a^2}\begin{pmatrix} 0 & -aE_1 & -aE_2 & -aE_3 \\ aE_1 & 0 & B_3 & -B_2 \\ aE_2 & -B_3 & 0 & B_1 \\ aE_1 & B_2 & -B_1 & 0 \end{pmatrix}. \tag{18}$$

The field equations (15), (16) and (17) rewritten in terms of $\vec{E}$ and $\vec{B}$ become

$$\vec{\nabla}\cdot\vec{E} = -\alpha\vec{\nabla}\phi\cdot\vec{B}. \tag{19a}$$

$$\partial_0(a^2\vec{E}) - \vec{\nabla}\times(a\vec{B}) = -\alpha[\partial_0\phi(a^2\vec{B}) - \vec{\nabla}\phi\times(a\vec{E})], \tag{19b}$$

$$\partial_0(a^2\vec{B}) - \vec{\nabla}\times(a\vec{E}) = 0, \tag{19c}$$

$$\vec{\nabla}\cdot\vec{B} = 0. \tag{19d}$$

For a spatially homogeneous pseudoscalar field equations (19a) and (19b) look like

$$\vec{\nabla}\cdot\vec{E} = 0 \tag{20a}$$

$$\partial_0(a^2\vec{E}) - \vec{\nabla}\times(a\vec{B}) = -\alpha\partial_0\phi(a^2\vec{B}). \tag{20b}$$

Taking the curl of (20b) and substituting into it the value of $\vec{E}$ from (19c) we obtain

$$\partial_0^2(a^2\vec{B}) + h(t)\partial_0(a^2\vec{B}) - \frac{\Delta^{(3)}(a^2\vec{B})}{a^2} - \frac{\alpha}{a}\partial_0\phi\vec{\nabla}\times(a^2\vec{B}) = 0, \tag{21}$$

where $\Delta^{(3)}$ stands for the three-dimensional Euclidean Laplacian operator.

Let us introduce

$$\vec{F}(\vec{x},t) \equiv a^2(t)\vec{B}(\vec{x},t) \tag{22}$$

and its Fourier transform

$$\vec{F}(\vec{k},t) = \frac{1}{(2\pi)^{3/2}}\int e^{-i\vec{k}\cdot\vec{x}}\vec{F}(\vec{x},t)d^3x. \tag{23}$$

Here the field $\vec{B}$ is an observable magnetic field entering into the expression for the Lorentz force. The field equation for $\vec{F}(\vec{k},t)$ is

$$\ddot{\vec{F}}(\vec{k},t) + h(t)\dot{\vec{F}}(\vec{k},t) + \left[\left(\frac{k}{a}\right)^2 - \frac{i\alpha}{a}\dot{\phi}\vec{k}\times\right]F(\vec{k},t) = 0, \tag{24}$$

where "dot" means the time derivative. This last equation can be further simplified: assuming $\vec{k} = (k, 0, 0)$ and defining the functions $F_\pm \equiv (F_2 \pm iF_3)/\sqrt{2}$ one arrives to

$$\ddot{F}_\pm + h\dot{F}_\pm + \left[\left(\frac{k}{a}\right)^2 \pm \alpha\frac{k}{a}\dot{\phi}\right] F_\pm = 0, \tag{25}$$

where we have omitted the arguments k and t.

Assuming that the electromagnetic field has a quantum origin (as all the fields in the cosmology of the early universe [14]) the modes of this field are represented by harmonic oscillators. Considering their vacuum fluctuations responsible for their birth we can neglect the small breakdown of the conformal symmetry and treat them as free. In conformal coordinates $(\eta, \vec{x})$ such that the Friedmann metric has the form

$$ds^2 = a^2(\eta)(d\eta^2 - dl^2), \tag{26}$$

the electromagnetic potential A_i with the gauge choice $A_0 = 0$, $\partial_j A^j = 0$ satisfies the standard harmonic oscillator equation of motion

$$\ddot{A}_i + k^2 A_i = 0. \tag{27}$$

Hence the initial amplitude of the field A_i behaves as $A_i = 1/\sqrt{2k}$, while the initial amplitude of the functions F is $\sqrt{k/2}$. The evolution of the field F during the inflationary period was described in [18], where it was shown that the growing solution at the end of inflation is amplified by some factor depending on the intensity of the interaction between the pseudoscalar field and magnetic field.

Here we are interested in the evolution of the magnetic field interacting with the pseudoscalar field after inflation, where our hypothetic Bang-to-Rip scenario takes place. More precisely, we would like to see how different types of scalar-pseudoscalar potentials and field dynamics providing the same cosmological evolution could be distinguished by their influence on the evolution of the magnetic field. We do not take into account the breaking of the conformal invariance during the inflationary stage and all the effects connected with this breakdown will be revealed only after the end of inflation and the beginning of the Bang-to-Rip evolution. This beginning is such that the value of the Hubble parameter, characterizing this evolution is equal to that of the inflation, i.e.

$$h(t_0) = \frac{t_R}{3t_0(t_R - t_0)} = h_{\text{inflation}} \simeq 10^{33}\text{s}^{-1}. \tag{28}$$

In turn, this implies that we begin evolution at the time moment of the order of 10^{-33}s.

We shall consider both the components F_+ and F_- and we shall dwell on the scenarios I and II described in the preceding section. Anyway, our assumption regarding the initial conditions for equation (25) can be easily modified in order to account for the previous possible amplification of primordial magnetic fields as was discussed by [18]. Thus, all estimates for the numerical values of the magnetic fields in today's universe should be multiplied by some factor corresponding to the amplification of the magnetic field during the inflationary stage. Hence, our results refer more to differences between various models of a post inflationary evolution than of the real present values of magnetic fields, whose amplification might be also combined effect of different mechanism [15, 17, 16, 18].

4. Generation of magnetic fields: numerical results

In this section we present the results of numerical simulations, for the two models I and II introduced in Sec. 2. In our models, the equation of motion for the modes $F_{\pm}(k,t)$ (25) reads¶:

$$\ddot{F}_{\pm}+\frac{t_R\,\dot{F}_{\pm}}{3t(t_R-t)}+\left[k^2\left(\frac{t_R-t}{t}\right)^{2/3}\pm\alpha\left(\frac{t_R-t}{t}\right)^{1/3}k\dot{\Phi}\right]F_{\pm}=0, \tag{29}$$

where Φ stands for the scalar field ϕ in the model I and for the phantom ξ in the model II, so that

$$\dot{\Phi}_I=\dot{\phi}=\frac{2}{3}\frac{\sqrt{t_R}}{t\sqrt{t_R-t}};\quad \dot{\Phi}_{II}=\dot{\xi}=\frac{2}{3}\frac{\sqrt{t_R}}{\sqrt{t}(t_R-t)}. \tag{30}$$

Equation (29) is solved for different values of the wave number k and the coupling parameter α. (The parameter α has the dimensionality inverse with respect to that of the scalar field; the wave number k has the dimensionality of inverse length; the time $t_R=10^{17}$s). Qualitatively we remark that in (25) the coupling term influence becomes negligible after some critical period. After that the magnetic fields in our different scenarios evolve as if the parameter α in (25) had been put equal to zero. Indeed, it can be easily seen that the interaction term vanishes with the growth of the cosmological radius a. Then the distinction between the two models is to be searched in the early time behavior of the field evolution.

Noting that in both our models I and II the time derivative $\dot{\Phi}$ is positive, by inspection of the linear term in equation (25) we expect the amplification to be mainly given for the mode F_- provided the positive sign for α is chosen; so we will restrict our attention on F_-. We can also argue that the relative strength of the last two terms in the left-hand side of (29) is crucial for determining the behavior of the solution: when the coupling term prevails (we remark that we are talking about F_- so this term is *negative* in our models) then we expect an amplification, while when the first term dominates we expect an oscillatory behavior. For future reference it is convenient to define

$$\mathbf{A}(t;\Phi)\equiv\frac{k}{a(t)\alpha\dot{\Phi}}=\frac{k}{\alpha\dot{\Phi}}\left(\frac{t_R-t}{t}\right)^{1/3}, \tag{31}$$

which is just the ratio between the last two terms in the left-hand side of equation (29).

Indeed our numerical simulations confirm these predictions. Let us consider the model I with $\alpha=1(t_P/m_P)^{1/2}$ and $k=10^{-55}l_P^{-1}$, where l_P is the Planck length. Such a value of the wave number k corresponds to the wave length of $1kpc$ at the present moment. We obtain an early-time amplification of about 2 orders of magnitude, with the subsequent oscillatory decay. Notice that the parameter $\mathbf{A}$ in this model at the beginning is very small: this corresponds to the dominance of the term proportional to $\dot{\Phi}$ and, hence, to the amplification of the field F_-. At the time scale of the order of $10^{51}t_P$, where t_P is the Planck time, this regime turns to that with big values of $\mathbf{A}$ where the influence of the term proportional to $\dot{\Phi}$ is negligible.

For the same choice of the parameters α and k in the model II the amplification is absent.

¶ The reader can easily verify that this equation is obtained imposing the normalization $a|_{\text{today}}=1$, where today-time is taken to be near the crossing of the phantom divide line, i.e. at $t\simeq t_R/2$. This implies in turn that at the beginning of the "Bang-to-Rip" evolution the cosmological radius is $a(t_0)\simeq10^{-17}$.



Figure 1. Plot of the ratio **A** for the model I, with the parameter choice $\alpha = 1(t_P/m_P)^{1/2}$, $k = 10^{-55} l_P^{-1}$. It can be easily seen that at a time scale of order $10^{51} t_P$ the ratio becomes greater than 1.



Figure 2. Plot of the time evolution of the absolute value of the complex field F (given in Planck units) in model I with the parameter choice $\alpha = 1(t_P/m_P)^{1/2}$, $k = 10^{-55} l_P^{-1}$. The behavior, as said above in the text, consists in an amplification till a time of order $10^{51} t_P$, after which the oscillations begin.



Figure 3. Plot of the time evolution of the absolute value of the complex field F (given in Planck units) in model I with the parameter choice $\alpha = 100(t_P/m_P)^{1/2}$, $k = 10^{-55} l_P^{-1}$. The behavior, consists in an amplification till a time of order $10^{54} t_P$, after which the oscillations begin.

In Figure 1 we present the time dependence of the function $\mathbf{A}$ for the model I for the values of $\alpha = 1(t_P/m_P)^{1/2}$ and $k = 10^{-55} l_P^{-1}$ chosen above. The Figure 2 manifests the amplification of the magnetic field in the model I.

Naturally the effect of amplification of the magnetic field grows with the coupling constant α and diminishes when the wave number k increases. In Figure 3 we display the results for the case of $\alpha = 100(t_P/m_P)^{1/2}$, which is admittedly extreme and possibly non realistic, but good for illustrative purposes. Here the amplification is more evident and extends for a longer time period.

Let us try to make some estimates of the cosmic magnetic fields in the universe today, using the correlation functions. The correlation function for the variable F is defined as the quantum vacuum average

$$G_{ij}(t, \vec{x} - \vec{y}) = \langle 0 | F_i(t, \vec{x}) F_j(t, \vec{y}) | 0 \rangle \tag{32}$$

and can be rewritten as

$$G_{ij}(t, \vec{x} - \vec{y}) = \int \frac{d^3k}{(2\pi)^3} e^{-i\vec{k}\cdot(\vec{x}-\vec{y})} F_i(t, \vec{k}) F_j^*(t, \vec{k}). \tag{33}$$

Integrating over the angles, we come to

$$G_{ij}(t, \vec{x} - \vec{y}) = \int \frac{k^2 dk}{2\pi^2} \frac{\sin(k|\vec{x} - \vec{y}|)}{k|\vec{x} - \vec{y}|} F_i(t, k) F_j^*(t, k). \tag{34}$$

To estimate the integral (34) we notice that the main contribution to it comes from the region where $k \approx 1/|\vec{x} - \vec{y}|$ (see e.g.[19]) and it is of order

$$\frac{1}{L^3} |F_i(t, 1/L)|^2, \tag{35}$$

where $L = |\vec{x} - \vec{y}|$. In this estimation the amplification factor is

$$\left| \frac{F(1/L)}{\sqrt{1/L}} \right|, \tag{36}$$

where the subscript i is not present since we have taken the trace over polarizations.

Now we are in a position to give numerical values for the magnetic fields at different scales in the model I for different values of the coupling parameter α. These values (see Table 1) correspond to three values of the coupling parameter α (1,10 and 100 $(t_P/m_P)^{1/2}$) [+] and to two spatial scales L determined by the values of the wave number k. We do not impose some physical restrictions on the value of α. It is easy to see that the increase of α implies the growth of the value of the magnetic field B. We also shall consider a rather large value of the wave number k corresponding to the physical wavelength $\sim 10^{-2}$ pc at the present moment when $a = 1$. While this scale looks too small for the description of large-scale cosmic magentic fields, we use it for illustrative purposes to underline the strong dependence of the amplification of magnetic field on the corresponing wavelength values.

Let us stress once again that we ignored the effects of amplification of the magnetic fields during inflation to focus on seizable effects during evolution.

Table 1. The Table displays the values of the magnetic field B corresponding to the chosen values of α and k. The length L refers to the present moment when $a = 1$.

	$\alpha = 1\sqrt{t_P/m_P}$	$\alpha = 10\sqrt{t_P/m_P}$	$\alpha = 100\sqrt{t_P/m_P}$
$k = 10^{-55} l_P^{-1}$ $(L = 1kpc)$	$B \sim 10^{-67} G$	$B \sim 10^{-60} G$	$B \sim 100 G$
$k = 10^{-50} l_P^{-1}$ $(L = 10^{-2} pc)$	$B \sim 10^{-55} G$	$B \sim 10^{-49} G$	$B \sim 10^{13} G$

Finally notice that our quantum "initial" conditions correspond to physical magnetic fields which for presented values of k are $B_{in} \sim k^2/a^2$ is equal to $10^{-34}G$ for $k = 10^{-55} l_P^{-1}$ and $B_{in} \sim 10^{-24}G$ for $k = 10^{-50} l_P^{-1}$.

5. Conclusion

We have seen that the evolution of the cosmic magnetic fields interacting with a pseudoscalar (pseudophantom) field is quite sensitive to the concrete form of the dynamics of this field in two-field models where different scalar field dynamics and potentials realize the same cosmological evolution.

We confirm the sensitivity of the evolution of the magnetic field with respect to its helicity given the sign of the coupling constant α and that the Φ is a monotonic function of time (as it is really so in our models). We give also some numerical estimates of the actual magnetic fields up to the factor of amplification of such fields during the inflationary period. The toy model of the Bang-to-Rip evolution studied in this paper, cannot be regarded as the only responsible for the amplification of cosmic magnetic fields implying their present observable values. It rather complements some other mechanisms acting before. However, the difference between cosmic magnetic

---

[+] It is useful to remark that at this length scales values of α less than 1 make the effect of the coupling of the magnetic field with the pseudoscalar field negligible.

fields arising in various models (giving the same expansion law after the inflation) is essential. It may provide a discriminating test for such models.

Naturally, the influence of the interaction between a pseudoscalar (phantom) field and a cosmic magnetic field on the dynamics of the latter depends on the velocity of change with time of the former. The larger is the time derivative of the pseudoscalar field, the more intensive is the growth of the magnetic field (remember that the evolution of the scalar field is monotonic). The results of numerical calculations illustrated in Sec. 4 confirm these qualitative considerations. Moreover, one can see that there exists a certain range of values of the wave number k (and hence of the corresponding wavelengths of the magnetic field) where the effect is stronger. Indeed, if k is too small the interaction term is small as well and the evolution of the magnetic field is damped. On the other hand if the wave number k is too large the interaction term is small compared to the oscillatory term, proportional as usual to k^2 and the evolution has practically oscillatory character. Thus, the study of interplay between the dynamics of global scalar fields providing the cosmological evolution and the magnetic fields looks promising. This interplay has another interesting aspect. The pseudoscalar-electromagnetic field interaction can imply a conversion of the photons into axions. Such an effect can cause the observable dimming of supenovae. While it was shown [20] that this effect cannot mimic the cosmic acceleration, it could nevertheless mimic the dark energy fluid with a phantom equation of state [21]. Thus, the interrelation amongst an electromagnetic field, a scalar and a phantom field can reveal some surprises.

Acknowledgments

This work was partially supported by Grants RFBR 08-02-00923 and LSS-4899.2008.2. The work of A.A. was supported by grants FPA2007-66665, 2005SGR00564, 2007PIV10046, by the Consolider-Ingenio 2010 Program CPAN (CSD2007- 00042) and Program RNP 2.1.1.1112.

References

[1] Riess A et al. 1998 *Astron. J.* **116** 1009; Perlmutter S J et al. 1999 *Astroph. J.* **517**, 565

[2] Sahni V and Starobinsky A A 2000 *Int. J. Mod. Phys.* D **9** 373; Padmananbhan T 2003 *Phys. Rep.* **380** 235; Peebles P J E and Ratra B 2003 *Rev. Mod. Phys.* **75** 559; Sahni V 2002 *Class. Quantum Grav.* **19** 3435; Copeland E J, Sami M and Tsujikawa S 2006 *Int. J. Mod. Phys.* D **15** 1753; Sahni V and Starobinsky A A 2006 *Int. J. Mod. Phys.* D **15** 2105; Frieman J A, Turner M S and Huterer D 2008 Dark energy and the accelerating universe *Preprint* 0803.0982 [astro-ph]

[3] Starobinsky A A 1998 *JETP Lett.* **68** 757; Boisseau B, Esposito-Farese G, Polarski D and Starobinsky A A 2000 *Phys. Rev. Lett.* **85** 2236; Burd A B and Barrow J D 1988 *Nucl. Phys.* B **308** 929; Barrow J D 1990 *Phys. Lett.* B **235** 40; Gorini V, Kamenshchik A, Moschella U and Pasquier V 2004 *Phys. Rev.* D **69** 123512; Zhuravlev V M, Chervon S V and Shchigolev V K 1998 *JETP* **87** 223; Chervon S V and Zhuravlev V M The cosmological model with an analytic exit from inflation *Preprint* gr-qc/9907051; Yurov A V Phantom scalar fields result in inflation rather than Big Rip *Preprint* astro-ph/0305019; Yurov A V and Vereshchagin S D 2004 *Theor. Math. Phys.* **139** 787; Guo Z K, Ohta N and Zhang Y Z 2007 *Mod. Phys. Lett.* A **22** 883; Guo Z K, Ohta N and Zhang Y Z 2005 *Phys. Rev.* D **72** 023504; Zhuk A 1996 *Class. Quant. Grav.* **13** 2163; Szydlowski M and Czaja W 2004 *Phys. Rev.* D **69** 083507; Szydlowski M and Czaja W 2004 *Phys. Rev.* D **69** 083518; Szydlowski M 2005 *Int. J. Mod. Phys.* A **20** 2443; Szydlowski M, Hrycyna O and Krawiec A 2007 *JCAP* **0706** 010; Vernov S Yu Construction of Exact Solutions in Two-Fields Models and the Crossing of the Cosmological Constant Barrier *Preprint* astro-ph/0612487

[4] Alam U, Sahni V, Saini T D and Starobinsky A A 2004 *Mon. Not. Roy. Astron. Soc.* **354** 275; Padmanabhan T and Choudhury T R 2003 *Mon. Not. Roy. Astron. Soc.* **344** 823; Choudhury T R and Padmanabhan T 2005 *Astron.Astrophys.* **429** 807; Alam U, Sahni V and Starobinsky A A 2004 *JCAP* **0406** 008; Wang Y and Freese K 2006 *Phys. Lett.* B **632**449; Upadhye A, Ishak M and Steinhardt P J 2005 *Phys. Rev.* D **72** 063501; Dicus D A and Repko W W 2004 *Phys. Rev.* D **70** 083527; Nesseris S and Perivolaropoulos L 2004 *Phys. Rev.* D **70** 043531; Lazkoz R, Nesseris S and Perivolaropoulos L 2005 *JCAP* **0511** 010

[5] Caldwell R R 2002 *Phys. Lett.* B **545** 23

[6] Alam U, Sahni V and Starobinsky A A 2007 *JCAP* **0702** 011; Nesseris S and Perivolaropoulos L 2007 *JCAP* **0702** 025

[7] Esposito-Farese G and Polarski D 2001 *Phys. Rev.* D **63** 063504; Vikman A 2005 *Phys. Rev.* D **71** 023515; Perivolaropoulos L 2005 *Phys. Rev.* D **71** 063503; McInnes B 2005 *Nucl. Phys.* B **718** 55; Aref'eva I Ya, Koshelev A S and Vernov S Yu 2005 *Phys. Rev.* D **72** 064017; Perivolaroupoulos L 2005 *JCAP* **0510** 001; Caldwell R R and Doran M 2005 *Phys. Rev.* D **72** 043527; Sahni V and Shtanov Yu 2003 *JCAP* **0311** 014; Sahni V and Wang L 2000 *Phys. Rev.* D **62** 103517

[8] Andrianov A A, Cannata F and Kamenshchik A Y 2005 *Phys. Rev.* D **72** 043531; Cannata F and Kamenshchik A Yu 2007 *Int. J. Mod. Phys.* D **16** 1683; Li M, Feng B and Zhang X 2005 *JCAP* **0512** 002

[9] Gannouji R, Polarski D, Ranquet A and Starobinsky A A 2006 *JCAP* **0609** 016

[10] Zhang X F, Li H, Piao Y S and Zhang X M 2006 *Mod. Phys. Lett.* A **21**231; Feng B, Wang X and Zhang X 2005 *Phys. Lett.* B **607** 35; Guo Z K, Piao Y S, Zhang X and Zhang Y Z 2005 *Phys. Lett.* B **608** 177; Hu W 2005 *Phys. Rev.* D **71** 047301; Perivolaropoulos L 2005 *Phys. Rev.* D **71** 063503; Caldwell R R and Doran M 2005 *Phys. Rev.* D **72** 043527.

[11] Andrianov A A, Cannata F and Kamenshchik A Yu 2006 *Int. J. Mod. Phys.* D **15** 1299; Andrianov A A, Cannata F and Kamenshchik A Yu 2006 *J. Phys.* A **39** 9975

[12] Andrianov A A, Cannata F, Kamenshchik A Yu and Regoli D 2008 *JCAP* **0802** 015

[13] Starobinsky A A 2000 *Grav. Cosmol.* **6** 157; Caldwell R R, Kamionkowski M and Weinberg N N 2003 *Phys. Rev. Lett.* **91** 071301

[14] Mukhanov V F and Chibisov G V 1981 *JETP Lett.* **33** 532

[15] Grasso D and Rubinstein H R 2001 *Phys. Rep.* **348** 163; Giovannini M 2004 *Int. J. Mod. Phys.* D **13** 391; Giovannini M 2008 *Lect.Notes Phys.* **737** 863

[16] Turner M S and Widrow L M 1988 *Phys. Rev.* D **77** 2743; Bamba K, Ohta N and Tsujikawa S, *Generic estimates for magnetic fields generated during inflation including Dirac-Born-Infeld theories*, 2008 *Preprint* 0805.3862 [astro-ph]; Diaz-Gil A, Garcia-Bellido J, Garcia Perez M and Gonzalez-Arroyo A, *Primordial magnetic fields from preheating at the electromagnetic scale*, 2008 *Preprint* 0805.4159 [hep-ph]

[17] Ratra B 1992 *Astrophys. J* **391** L1; Garretson W D, Field G B and Carroll S M 1992 *Phys. Rev.* D **46** 5346; Dolgov A D 1993 *Phys. Rev.* D **48** 2499; Field G B and Carroll S M 2000 *Phys. Rev.* D **62** 103008; Finelli F and Gruppuso A 2001 *Phys. Lett.* B **502** 216; Bamba K and Sasaki M 2007 *JCAP* **0702** 030

[18] Anber M M, Sorbo L *JCAP* 2006 **0610** 018

[19] Mukhanov V 2005 *Physical Foundations of Cosmology* (Cambridge University Press)

[20] Mirizzi A, Raffelt G G and Serpico P D 2008 *Lect. Notes Phys.* **741** 115

[21] Csaki C, Kaloper N and Terning J 2005 *Annals of Phys* **317** 410